

CONTINUITY OF THE ITÔ-MAP FOR HÖLDER ROUGH PATHS WITH APPLICATIONS TO THE SUPPORT THEOREM IN HÖLDER NORM

PETER K. FRIZ

ABSTRACT. Rough Path theory is currently formulated in p -variation topology. We show that in the context of Brownian motion, enhanced to a Rough Path, a more natural Hölder metric ρ can be used. Based on fine-estimates in Lyons' celebrated Universal Limit Theorem we obtain Lipschitz-continuity of the Itô-map (between Rough Path spaces equipped with ρ). We then consider a number of approximations to Brownian Rough Paths and establish their convergence w.r.t. ρ . In combination with our Hölder ULT this allows sharper results than the p -variation theory. Also, our formulation avoids the so-called control functions and may be easier to use for non Rough Path specialists. As concrete application, we combine our results with ideas from [MS] and [LQZ] and obtain the Stroock-Varadhan Support Theorem in Hölder topology as immediate corollary.

1. INTRODUCTION

1.1. Background in Rough Path theory. Over the last years T.Lyons and co-authors developed a general theory of integration and differential equations of form

$$(1.1) \quad dy_t = f(y_t)dx_t.$$

To include the important example of stochastic differential equations, x must to be allowed to be "rough" in some sense. Standard Hölder-regularity of Brownian motion, for instance, implies finite p -variation only for $p > 2$. Another issue is to explain (deterministically) the difference between SDEs based on Stratonovich- vs.

Date: February 1, 2008.

Key words and phrases. Rough Path theory, Itô-map, Universal Limit Theorem, p -variation vs. Hölder regularity, Support Theorem.

Itô-integrals. Last not least, motivated from examples like Fractional Brownian motion, driving signals much rougher than Brownian motion should be included.

All this has been accomplished in a very satisfying way and the reader can nowadays find the general theory exposed in [L98], [LQ], [Le].

Loosely speaking, for general $p \geq 1$, one needs to "enhance" the driving signal x , with values in some Banach-space V , to $X \in V \oplus V^{\otimes 2} \dots \oplus V^{\otimes [p]}$ such that the resulting object X satisfies certain algebraic¹ and analytic conditions. For x of finite variation, this enhancement will simply consist of all the iterated integrals of x ,

$$X_{s,t}^k := \int_{s < u_1 < \dots < u_k < t} dx_{u_1} \otimes \dots \otimes dx_{u_k}, \quad k = 1, \dots, [p].$$

These are the *Smooth Rough Paths*. Consider a time-horizon of $[0, 1]$ (valid for the rest of the paper) and introduce the p -variation metric, defined as

$$d(X, Y) = \max_{k=1, \dots, [p]} \left(\sup_D \sum_l |X_{t_{l-1}, t_l}^k - Y_{t_{l-1}, t_l}^k|^{p/k} \right)^{k/p},$$

where $\sup_D$ runs over all finite divisions of $[0, 1]$. Here $|\cdot|$ denotes (compatible) tensor norms in $V^{\otimes k}$. Closure of Smooth Rough Paths w.r.t. this metric yields the class of *Geometric Rough Paths*², denoted by $G\Omega_p(V)$. The solution-map, also called *Itô-map*, to (1.1) is then a continuous map from $G\Omega_p(V) \rightarrow G\Omega_p(W)$, provided $f : W \rightarrow L(V, W)$ satisfies mild regularity conditions. This is Lyons' celebrated *Universal Limit Theorem*. In particular, smooth approximations $X(n)$ which converge in p -variation to $X \in G\Omega_p(V)$ will cause the corresponding solutions $Y(n)$ to converge to Y in p -variation. Hence, one deals with some kind of generalized Stratonovich theory.

However, the so important case of $p \in (2, 3)$, on which this paper will focus, allows for more. Following [LQ] p149 and also [L98] the driving signal only needs to be a *Rough Path of finite p -variation*. By definition, this is a continuous map

$$(s, t) \rightarrow (X_{s,t}^1, X_{s,t}^2) \in V \oplus V^{\otimes 2},$$

¹For *algebraic* convenience X is often enhanced to $\mathbb{R} \oplus V \oplus V^{\otimes 2} \dots \oplus V^{\otimes [p]}$ with scalar component constant 1.

²A slightly weaker definition of Geometric Rough Path appeared in [L98]. We follow the more recent [LQ].

where $0 \leq s \leq t \leq 1$, satisfying the algebraic condition

$$(1.2) \quad X_{s,u}^1 = X_{s,t}^1 + X_{t,u}^1 \quad \text{and} \quad X_{s,u}^2 = X_{s,t}^2 + X_{t,u}^2 + X_{s,t}^1 \otimes X_{t,u}^1,$$

whenever $s \leq t \leq u$, and the analytic condition $d(X, 0) < \infty$ i.e.

$$(1.3) \quad \sup_D \sum_l |X_{t_{l-1}, t_l}^k|^{p/k} < \infty, \quad k = 1, 2 \quad .$$

(Sometimes we refer to $k = 1, 2$ as *first* resp. *second level*.) The class of Rough Paths of finite p -variation is denoted by $\Omega_p(V)$. Clearly,

$$\{\text{Smooth Rough Paths}\} \subset G\Omega_p(V) \subset \Omega_p(V).$$

Condition (1.2) is known as *Chen relation* and expresses simple additive properties whenever X^2 is obtained as *some* iterated integral including the cases of Stratonovich- resp. Itô-Enhanced Brownian Motion (EBM), where

$$X_{s,t}^1 = \beta_{s,t}^1 = \omega_t - \omega_s,$$

the increments of Brownian Motion on $V = \mathbb{R}^d$, defined on the usual Wiener-space $(C([0, 1], \mathbb{R}^d), \mu)$. $X_{s,t}^2 = \beta_{s,t}^2$ is defined via stochastic integration,

$$(1.4) \quad \int_s^t (\omega_r^i - \omega_s^i) \circ d\omega_r^j \quad \text{resp.} \quad \int_s^t (\omega_r^i - \omega_s^i) d\omega_r^j.$$

Clearly, there is a modification of β^2 , denoted by the same name, such that β^2 (and hence β) is continuous in s, t for all ω . With this choice, it is well-known that (both Stratonovich- and Itô-)EBM are a.s. elements of $\Omega_p(V)$, also called *Brownian Rough Paths*. Indeed, this will follow from a.e. Stratonovich-EBM $\beta = \beta(\omega) = (\beta^1, \beta^2)$ being a Geometric Rough Path and this is the (somewhat computational) approach in [LQ], [Le].

We emphasize once more, that for $p \in (2, 3)$ and driving signal in $\Omega_p(V)$ a complete theory is available, covering both Itô- and Stratonovich-SDEs. In particular, the Universal Limit Theorem holds, see [LQ], p164 and Section 4.

1.2. Definitions and outline. Our focus being on Brownian Rough Paths, let $p \in (2, 3)$ and $V = \mathbb{R}^d$ from here on. Given a Rough Path

$$(1.5) \quad \|X\| := \max_{k=1,2} \sup_{0 \leq s < t \leq 1} \frac{|X_{s,t}^k|}{|t-s|^{k/p}}.$$

We call Rough Paths with $\|X\| < \infty$ *Hölder Rough Paths* and write $X \in H\Omega_p(V)$. This condition is similar to (1.3) but stronger. Note $H\Omega_p(V) \subset \Omega_p(V)$. Clearly, $H\Omega_p(V)$ becomes a metric space under $\rho(X, Y) := \|X - Y\|$. (The norm $\|\cdot\|$ appeared in [L95] but seems to have disappeared in the current *p-variation* Rough Path theory.) In **Section 2** we use Kolmogorov's criterion to prove a 2-parameter Hölder-regularity of an object intimately related to β . This implies $\beta \in H\Omega_p(V)$. (Such a result was mentioned without proof in [L95] and might be contained in unpublished thesis-work, [S].) As further corollary, we have the well-known $\beta \in \Omega_p(V)$. In **Section 3**, we establish convergence in ρ -metric of piecewise linear approximations based on nested partitions. This gives a simple and novel proof that Stratonovich-EBM is an element of $G\Omega_p(V)$. (Our proof combines the preceding regularity result with soft martingale and compactness arguments. In a sense, all dyadic approximations done by hand in [LQ], [Le] are isolated in our initial application of Kolmogorov's criterion.) At last, the results are extended to adapted dyadics approximations.

Section 4 is a recall of Lyons' Universal Limit Theorem. In **Section 5** we translate his fine estimates in terms of *control functions* to plain Lipschitz-continuity of the Itô-map from $(H\Omega_p(V), \rho)$ to $(H\Omega_p(W), \rho)$.

Finally, in **Section 6** we apply our refined Rough Path machinery to prove the celebrated Support Theorem in Hölder-topology. As observed in [MS], the proof can be reduced to two convergence results (one for each inclusion) and these follow immediately from our results.

Remarks: -The use of Rough Path theory to prove the Support Theorem was first carried out in [LQZ]. Using the standard (= *p-variation*) Rough Path machinery the well-known Hölder-topology result ([BGL], [MS], [ST]) was not recovered. We also note, that the approach in [LQZ] relies on correlation inequalities.

-A recent preprint, [Gu], proposes to re-prove rough path theory in a pure Hölder context, apparently motivated by (1.5).

-Forthcoming joint-work with Nicolas Victoir will (among other things) contain a rough paths proof of the support theorem in yet a stronger norm (modulus norm).³

2. HÖLDER-REGULARITY OF ENHANCED BROWNIAN MOTION

As before, let $(\beta^1, \beta^2) = \beta = \beta(\omega)$ be (Itô -or Stratonovich) EBM based on d -dimensional BM, chosen s.t. $(s, t) \mapsto \beta_{s,t}(\omega)$ is continuous for all ω . Recall $0 \leq s, t \leq 1$, $p \in (2, 3)$.

Theorem 2.1. *Let $\gamma \in (0, 1/2 - 1/p)$. Introduce the following 2-parameter processes*

$$(2.1) \quad Z_{s,t}^k = \frac{\beta_{s,t}^k}{|t-s|^{k/p}} \in V^{\otimes k} \cong \mathbb{R}^{d^k} \quad k = 1, 2,$$

whenever $t > s$ and set them zero otherwise. Then for μ -a.e. ω $Z^k = Z^k(\omega)$, as function of s, t , is Hölder continuous of any exponent $k\gamma$. Also,

$$(2.2) \quad \sup_{s \neq s', t \neq t'} \frac{|Z_{s,t}^k - Z_{s',t'}^k|}{[|t' - t| + |s' - s|]^{k\gamma}} \in L^{\infty-} := \cap_{q < \infty} L^q.$$

The following lemma is proved in the appendix.

Lemma 2.2. *Fix $i, j \in \{1, \dots, d\}$ and set $Z = Z^{2,ij}$. Then there is $c = c(p)$ s.t.*

$$E[|Z_{s,t} - Z_{s',t'}|^2] \leq c[|s' - s| + |t' - t|]^{2(1-2/p)}.$$

Proof of Theorem 2.1.

We only prove $k = 2$. The first level i.e. $k = 1$ is similar but easier. As before, fix i, j and set $Z_{s,t} = Z_{s,t}^{k,ij}$. By equivalence of finite dimensional norms, it suffices to show that Z is Hölder. We first consider Itô-EBM. With this choice $Z_{s,t}$ is an element of the second Wiener-Itô-chaos and a moment estimate similar to the

³The overlap to [FV] is kept to a minimum. In particular, 2-parameter regularity and convergence, definition (1.5) and Lipschitz-continuity of Itô-map are not contained in [FV].

Gaussian case is available ([RY], p207). With lemma 2.2,

$$\begin{aligned} E[|Z_{s,t} - Z_{s',t'}|^q] &\leq c(q) \left(E[|Z_{s,t} - Z_{s',t'}|^2] \right)^{q/2} \\ &\leq c[|s' - s| + |t' - t|]^{(1-2/p)q}. \end{aligned}$$

We can choose q arbitrarily large and, by Kolmogorov's criterion as found in [RY], obtain Hölder regularity for any exponent γ less than $1 - 2/p$.

The only thing left to consider is $Z = Z^{2,ij}$ based on Stratonovich-EBM. Only on the diagonal $i = j$ there is a non-zero difference (coming from the quadratic variation of BM),

$$Z_{s,t}^{\text{Itô}} - Z_{s,t}^{\text{Stratonovich}} = \frac{1}{2}|t - s|^{1-2/p}.$$

As composition of the Lipschitz-map $(s, t) \rightarrow |t - s|$ and a $(1 - 2/p)$ -Hölder map, the map $(s, t) \mapsto |t - s|^{1-2/p}$ is itself $(1 - 2/p)$ -Hölder. This implies the γ -Hölder-regularity of $Z = Z^{\text{Stratonovich}}$. The proof is finished. QED

Corollary 2.3. *Enhanced Brownian motion β is μ -a.s. an element of $H\Omega_p(V) \subset \Omega_p(V)$. More precisely, there exists $C = C(\omega) \in L^\infty$ such that for all $s \leq t$ and $k = 1, 2$*

$$(2.3) \quad |\beta_{s,t}^k|^{p/k} \leq C|t - s|.$$

(In Lyons' terminology, β admits the additive control $C|t - s|$.)

Proof of Corollary 2.3.

The continuous function $(s, t) \mapsto Z_{s,t}^k$ achieves its maximum $C(\omega)$ which is easily estimated by the (2.2). QED

Remark: We showed that a.s. Z^k is an element of the Hölder-space $C^{0,k\gamma}([0, 1]^2, V^{\otimes k})$. Moreover, (2.2) can be used as a norm.

3. APPROXIMATIONS TO BROWNIAN ROUGH PATHS

3.1. Piecewise linear nested approximations. Based on piecewise linear nested approximations $\omega(n)$ of the underlying Brownian path ω we construct a Smooth Rough Path denoted by $\beta(n) = (\beta^1(n), \beta^2(n))$. We assume that these partitions are deterministically chosen and that their mesh goes to 0. Note that the commonly used piecewise linear *dyadic* approximations (see [ST], [M], [LQ], ...) fall into the considered class.

As before, introduce Z^1, Z^2 based on Stratonovich-EBM β and, similarly $Z^1(n), Z^2(n)$ based on $\beta(n)$. The matrix-valued processes $Z^2, Z^2(n), \dots$ split into symmetric and anti-symmetric parts. For instance,

$$Z^2 = \hat{Z}^2 + A^2,$$

where $\hat{\cdot}$ indicates symmetrization.

Theorem 3.1. *Let $p \in (2, 3), \gamma \in (0, 1/2 - 1/p)$. For first level and symmetric part of the second level, convergence of the approximations holds a.s. in Hölder-space of exponent γ ,*

$$(3.1) \quad \|Z^1 - Z^1(n)\|_{C^{0,\gamma}} \rightarrow 0,$$

$$(3.2) \quad \|\hat{Z}^2 - \hat{Z}^2(n)\|_{C^{0,\gamma}} \rightarrow 0.$$

For the antisymmetric part A^2 , based on the Lévy-area of the the underlying BM, we have a.s.

$$(3.3) \quad \|A^2 - A^2(n)\|_{C^{0,2\gamma}} \rightarrow 0.$$

Remark: Given that Z^2 is itself 2γ -Hölder it is quite possible that $Z^2(n) \rightarrow Z^2$ in $C^{0,2\gamma}$, but none of our conclusions will rely on this.

Corollary 3.2. *There exists a random constant $C < \infty$ a.s. and a random sequence $a_n(\omega) \rightarrow 0$ a.s. such that, for $s \leq t, k = 1, 2$*

$$(3.4) \quad |\beta_{s,t}^k(n)|, |\beta_{s,t}^k| \leq [C(\omega)|t - s|]^{k/p}$$

and

$$(3.5) \quad |\beta_{s,t}^k - \beta_{s,t}^k(n)| \leq a_n(\omega)|t - s|^{k/p}.$$

(In Lyons' terminology, the additive control $C(\omega)|t - s|$ is uniform for the entire sequence $\beta(n)$ and controls the convergence. ⁴)

These estimates translate to

Corollary 3.3. *The (Smooth Rough Path-) approximations $\beta(n)$ converge to β in Hölder-metric ρ ,*

$$\rho(\beta(n), \beta) \rightarrow 0 \quad \text{a.s.}$$

(Since this implies convergence in p -variation metric d we identify, en passant, Stratonovich-EBM β as a Geometric Rough Path.)

⁴Strictly speaking, $(C \vee 1)|t - s|$ will be the required control.

Proof of Theorem 3.1. We are able to do most of the work for levels $k = 1, 2$ at the same time. For the moment, fix $i \neq j$. For $k = 1$ set

$$Z = Z^{1,i},$$

while for $k = 2$ set

$$Z = Z^{2,ij}.$$

Either way, we have a real-valued 2-parameter process, $k\gamma$ -Hölder according to Theorem 2.1. This means for some L^∞ -r.v. L we have the inequality

$$-L[|t' - t| + |s' - s|]^{k\gamma} \leq Z_{s,t} - Z_{s',t'} \leq L[|t' - t| + |s' - s|]^{k\gamma}.$$

Now condition w.r.t. $\mathcal{F}_n := \sigma(\beta_{k/2^n} : k = 0, \dots, 2^n)$. Set $L_n = E[L|\mathcal{F}_n]$. It is not hard to see ([M], p216) that $E[Z|\mathcal{F}_n] = Z(n)$. (In the case $k = 2$, here is where we use $i \neq j$.) Hence,

$$-L_n[|t' - t| + |s' - s|]^{k\gamma} \leq Z_{s,t}(n) - Z_{s',t'}(n) \leq L_n[|t' - t| + |s' - s|]^{k\gamma}.$$

Note that L_n is an L^∞ -bounded martingale. By Doob's L^p -inequality we see that,

$$C(\omega) := \sup_n L_n(\omega)$$

is also in L^∞ . Consequently,

$$\sup_{s,t,s',t'} \frac{|Z_{s,t}(n) - Z_{s',t'}(n)|}{[|t' - t| + |s' - s|]^{k\gamma}} \leq L_n \leq C.$$

This implies that the sequence $(Z(n))$ is bounded in the Hölder-space $C^{0,k\gamma}$. We could have started with $\gamma + \epsilon$ as long as $\gamma + \epsilon < 1/2 - 1/p$. Then the conclusion is boundedness in $C^{0,k(\gamma+\epsilon)}$ and by compactness there is a convergent subsequence in $C^{0,k\gamma}$. But every possible limit point is identified as Z itself, since for s, t fixed, $Z_{s,t}(n) \rightarrow Z_{s,t}$ by martingale convergence. This implies that $Z(n)$ actually converges to Z in $C^{0,k\gamma}$ and the same holds true for the antisymmetric part of $Z_{s,t}^2$ itself since all the diagonal, $i = j$, is zero.

At last, we need to consider the case

$$Y_{s,t} := 2Z_{s,t}^{2,ii} = \frac{(\beta_t^i - \beta_s^i)^2}{|t - s|^{2/p}} = (Z_{s,t}^{1,i})^2 =: (Z_{s,t})^2.$$

Similarly define $Y(n), Z(n)$. We claim that $Y(n)$ tends to Y in $C^{0,\gamma}$. Certainly, for s, t fixed $Y_{s,t}(n) \rightarrow Y_{s,t}$ which identifies every possible

limit point of $Y(n)$. Hence it suffices, by the same compactness argument as before, to show that $Y(n)$ is uniformly bounded in $C^{0,\gamma}$. But $Y(n) = [Z(n)]^2$ with $Z(n) = Z^{1,i}(n)$ and this last sequence was shown earlier to be uniformly bounded (even convergent) in $C^{0,\gamma}$. On the other hand, the map $x \mapsto x^2$ is (locally) Lipschitz, and since $\{Z(n)\}$ remains in a ball in $C^{0,\gamma}$ we conclude that $\{Y(n)\}$ remains in a (possibly larger) ball in $C^{0,\gamma}$ as well. The claim is proved. Together with the earlier results for $k = 2, i \neq j$ we find that $\hat{Z}^2(n)$, the symmetric part of $Z^2(n)$, converges in $C^{0,\gamma}$ to $\hat{Z}$. The proof is finished. QED

Proof of Corollary 3.3.

For $k = 1$ the estimates (3.4) and (3.5) are an immediate consequence of (3.1) and $Z^1 \in C^{0,\gamma}$.

Towards $k = 2$ we first consider the anti-symmetric part of $\beta_{s,t}^2 - \beta_{s,t}^2(n)$. But this is simply bounded by

$$a_1(n, \omega) |t - s|^{2/p}$$

where $a_1(n)$ denotes the l.h.s. of (3.3). Towards the symmetric part of $\beta_{s,t}^2 - \beta_{s,t}^2(n)$ observe that

$$\hat{\beta}_{s,t}^{2,ij} = \frac{1}{2} \beta_{s,t}^{1,i} \beta_{s,t}^{1,j}.$$

It suffices to estimate one component of $\hat{\beta}^2 - \hat{\beta}^2(n)$, namely,

$$\begin{aligned} |\beta_{s,t}^{1,i} \beta_{s,t}^{1,j} - \beta_{s,t}^{1,i}(n) \beta_{s,t}^{1,j}(n)| &\leq |\beta_{s,t}^{1,i}| |\beta_{s,t}^{1,j} - \beta_{s,t}^{1,j}(n)| \\ &\quad + |\beta_{s,t}^{1,i} - \beta_{s,t}^{1,i}(n)| |\beta_{s,t}^{1,j}(n)| \end{aligned}$$

for arbitrary i, j . From (3.1) it follows that

$$|\beta_{s,t}^1 - \beta_{s,t}^1(n)| \leq a_2(n, \omega) |t - s|^{1/p},$$

where $a_2(n)$ denotes the l.h.s. of (3.1). Together with the uniform estimates (3.4) we conclude that

$$|\hat{\beta}_{s,t}^2 - \hat{\beta}_{s,t}^2(n)| \leq a_3(n, \omega) |t - s|^{2/p},$$

where a_3 is a deterministic constant times $C(\omega)a_2$. For

$$a(n) := \max\{a_1(n), a_3(n)\}$$

estimate (3.5) will then hold true. Also, (3.4) follows by the triangle-inequality and the regularity of β^2 . QED.

3.2. Adapted dyadic approximations. Now let $\omega(n)$ be the dyadic piecewise linear approximation to a Brownian path ω , i.e. piecewise linear from $\omega_{i/2^n}$ to $\omega_{(i+1)/2^n}$. Note, that $\omega(n)$ is not adapted to the Brownian filtration. This suggests to look at the following *adapted* approximation,

$$\omega_t^{ad}(n) = \omega_{(t-2^{-n}) \vee 0}(n).$$

We can lift the path $\omega^{ad}(n)$ to make it a (Smooth) Rough Path, which we denote by $\beta^{ad}(n)$. Similarly, and as before, $\omega(n)$ is lifted to $\beta(n)$.

Corollary 3.4. *The Smooth Rough Paths $\beta^{ad}(n)$ converge a.s. to Stratonovich-EBM β in Hölder-metric ρ (and consequently in p -variation metric d).*

Proof: Introduce a shift-operator on the path-level s.t. for any path $x(t) \in V$,

$$\tau^\epsilon : x(\cdot) \mapsto x((\cdot - \epsilon) \vee 0).$$

This lifts to a map on Rough Paths. With $\epsilon = 1/2^n$, Then,

$$\beta^{ad}(n) = \tau^\epsilon(\beta(n)).$$

In particular, for the second level,

$$\beta_{s,t}^{ad,2}(n) = \beta_{(s-\epsilon) \vee 0, (t-\epsilon) \vee 0}^2.$$

By the triangle inequality,

$$\rho(\tau^\epsilon(\beta(n)), \beta) \leq \rho(\tau^\epsilon(\beta(n)), \tau^\epsilon(\beta)) + \rho(\tau^\epsilon(\beta), \beta).$$

Since $\rho(\beta(n), \beta) \rightarrow 0$ so does the first term on the r.h.s. (ρ is insensitive to shift). So all that remains to show is that $\rho(\tau^\epsilon(\beta), \beta) \rightarrow 0$ as ϵ tends to zero. Written out, this means

$$\max_{k=1,2} \sup_{0 \leq s < t \leq 1} \frac{|\beta_{(s-\epsilon) \vee 0, (t-\epsilon) \vee 0}^k - \beta_{s,t}^k|}{|t-s|^{k/p}}$$

goes to zero with ϵ . We can estimate this by

$$\sup_{0 \leq s < t \leq \epsilon} (...) + \sup_{0 \leq s < \epsilon \leq t \leq 1} (...) + \sup_{\epsilon \leq s < t \leq 1} (...).$$

Each part is easily seen to converge to 0 with ϵ by using the (s, t) Hölder-property of $Z_{s,t}^k = \beta_{s,t}^k / |t-s|^{k/p}$ established in Theorem 3.1. QED

Clearly, the last corollary implies that on path level and in Hölder-norm with exponent less than $1/p$,

$$\omega^{ad}(n) \rightarrow \omega.$$

Then, trivially,

$$T_n(\omega) = \omega - \omega^{ad}(n) + h \rightarrow h,$$

for, say, any piecewise linear dyadic path h . We will lift this convergence result to Rough Path level. As in [LQZ] we shall denote the lifts of dyadic piecewise linear paths by $\mathcal{D}$, a set of smooth rough paths. Before doing so, recall the Stratonovich enhancement of Brownian motion,

$$\omega \rightarrow \beta_{s,t}(\omega) = (\omega_t - \omega_s, \frac{1}{2}(\omega_t - \omega_s)^{\otimes 2} + A_{s,t}) \in V \oplus V^{\otimes 2}.$$

Due to the Lévy-area $A_{s,t} = A_{s,t}(\omega)$ this is only an a.s. defined function of ω (although we picked a modification, determined up to indistinguishability, s.t. $(s, t) \rightarrow \beta_{s,t}(\omega)$ is continuous). We saw in section 2 that $\beta \in H\Omega_p(V)$ for μ -a.e. ω . By Girsanov's theorem, $\beta(T_n(\omega))$ is well-defined and in $H\Omega_p(V)$ for μ -a.e. ω .

Corollary 3.5. *The Hölder Rough Paths $\beta(T_n^h(\omega))$ converge a.s. in Hölder-metric p to the (Smooth) Rough Path $(h^1, h^2) \in \mathcal{D}$, associated to the piecewise linear dyadic path $h(\cdot)$.*

Proof: It suffices to consider the Lévy-area, more specifically off-diagonal term of the 2nd-level. That is, we want to show that, for $i \neq j$,

$$\sup_{s < t} \frac{|\beta_{s,t}^{2,ij}(T_n^h(\omega)) - h_{s,t}^{2,ij}|}{|t - s|^{2/p}}$$

tends to zero. To this end, a Riemann-sum approximation shows that the following expansion holds a.s. (we omit s, t , the following integrations are understood over the simplex $\{(u_1, u_2) : s \leq u_1 \leq u_2 \leq t\}$).

$$\begin{aligned} \beta^{2,ij}(T_n^h(\omega)) &= \beta^{2,ij}(\omega - \omega^{ad}(n) + h) = h^{2,ij} \\ &+ \beta^{2,ij}(\omega) + \int d\omega^{ad,i}(n) d\omega^{ad,j}(n) - \int d\omega^i d\omega^{ad,j}(n) - \int d\omega^{ad,i}(n) d\omega^j \\ &+ \int dh^i d\omega^j - \int dh^i d\omega^{ad,j}(n) + \int d\omega^i dh^j - \int d\omega^{ad,i}(n) dh^j. \end{aligned}$$

(All iterated integrals here make sense as Young-integrals.) Note that the last corollary implies that

$$(3.6) \quad \int d\omega^{ad,i}(n)d\omega^{ad,j}(n) \rightarrow \beta^{2,ij}(\omega)$$

w.r.t. the second level part of the metric ρ (just keep $k = 2$ in its definition). Clearly, $\int d\omega^i d\omega^{ad,j}(n)$ and $\int d\omega^{ad,i}(n)d\omega^j$ are even better approximations to $\beta^{2,ij}(\omega)$ and hence converge to the latter.

⁵ Then all four terms together in the second line (r.h.s.) above will converge to zero.

As for the third line, observe that (3.6) holds for a.e. Brownian path ω . By Girsanov's theorem we have a.s. convergence after replacing ω by $\omega + h$. Now use an expansion as before,

$$\beta^{2,ij}(\omega + h) = \beta^{2,ij}(\omega) + h^{2,ij} + \int dh^i d\omega^j + \int d\omega^i dh^j.$$

Similarly, expand

$$\int d(\omega^{ad,i}(n) + h^i(n))d(\omega^{ad,j}(n) + h^j(n)).$$

(Note that $h(n) = h \in \mathcal{D}$ for all n large enough.) As already mentioned, (3.6) still holds after replacing ω by $\omega + h$. Together with the expansions, this gives exactly the required cancelation (as $n \rightarrow \infty$) of the third line above. QED

Remark: The preceding proof involves a perturbation of the rough path β (essentially) in a Cameron-Martin-type direction, (h^1, h^2) . Such and more general perturbations have been studied systematically in [LQ97], see also [LQ]. Indeed, we could have based our proof on some of their general results.

4. A PRIMER ON THE UNIVERSAL LIMIT THEOREM

We just summarize and plug together a few statements from [LQ]. Set $V = \mathbb{R}^d$, $W = \mathbb{R}^N$. As before, $p \in (2, 3)$.

Recall that a *control function*⁶ is, by definition, a non-negative continuous function w on $\{0 \leq s \leq t \leq 1\}$, super-additive, that is, $w(s, t) + w(t, u) \leq w(s, u)$ and hence zero on the diagonal.

⁵This is easily made precise by partial conditioning in the proof of Theorem 3.1, that is condition w.r.t. the i th resp. j th component only.

⁶The reader who does not want to know about control functions may jump to the statement of Theorem 5.1 directly.

Theorem 4.1. *Let $f \in C^3(W, L(V, W))$, globally Lipschitz. Then the Itô-map from $\Omega_p(V) \rightarrow \Omega_p(W)$, obtained from “enhancing”*

$$dy = f(y)dx, y(0) = y_0,$$

is continuous w.r.t. the p -variation metric. Moreover, if $w = w(s, t)$ is a control function such that (always for all s, t)

$$(4.1) \quad |X_{s,t}^k|, |\hat{X}_{s,t}^k| \leq w(s, t)^{k/p}, \quad k = 1, 2$$

and

$$(4.2) \quad |X_{s,t}^k - \hat{X}_{s,t}^k| \leq \epsilon w(s, t)^{k/p}, \quad k = 1, 2,$$

then there exists a constant $C = C(\max w) = C(\max w, f, p, y_0)$ such that

$$|Y_{s,t}^k|, |\hat{Y}_{s,t}^k| \leq (Cw(s, t))^{k/p}, \quad k = 1, 2$$

and

$$|Y_{s,t}^k - \hat{Y}_{s,t}^k| \leq \epsilon (Cw(s, t))^{k/p}, \quad k = 1, 2.$$

Proof: The fine estimate in terms of control functions w are stated and proved in [LQ], p163. For the readers convenience, let us quickly show how to obtain continuity w.r.t p -variation metric $d(., .)$. Assume $d(X(n), X) \rightarrow 0$. Clearly, for any subsequence, $d(X(n_i), X) \rightarrow 0$. By ⁷ [LQ], p51, there exists a further subsequence $(n_{i_j} =: n_j)$ and a control function w such that

$$|X_{s,t}^k|, |X_{s,t}^k(n_j)| \leq w(s, t)^{k/p}, \quad k = 1, 2$$

for any j and

$$|X_{s,t}^k - X_{s,t}^k(n_j)| \leq 2^{-j} w(s, t)^{k/p}, \quad k = 1, 2.$$

Hence, by the ULT estimates above,

$$|Y_{s,t}^k - Y_{s,t}^k(n_j)| \leq 2^{-j} Cw(s, t)^{k/p}, \quad k = 1, 2.$$

Using the super-additivity this implies $d(Y(n_j), Y) \rightarrow 0$. Since we were able to extract, from any subsequence, a further convergent subsequence with limit Y , it is clear that $d(Y(n), Y) \rightarrow 0$. Hence $X \rightarrow Y$ is continuous in p -variation metric. QED.

⁷Unfortunately, the cited page contains a misprint. The supremum of superadditive functions may loose this property. However, all this is readily fixed and since we will not rely on the results we leave the corrections to the reader.

Remarks: (1) The fine-estimates exhibit some kind of Lipschitz behavior which is not visible in terms of p -variation metric.
 (2) Since a.s. $d(\beta(n), \beta) \rightarrow 0$ Theorem (4.1) applies and yields a Wong-Zakai-type result. However, constructing a control w as above is certainly a bad idea, since Corollary 3.2 provides us with a much easier control, which indeed controls the whole sequence rather than just a subsequence. Essentially, $w(s, t) = c|t - s|$ for some (random) constant c . We will now exploit these observations.

5. LIPSCHITZ REGULARITY OF THE ITÔ-MAP FOR HÖLDER ROUGH PATHS

Theorem 5.1. *Under assumptions on f as in Theorem 4.1 the Itô-map $F : X \rightarrow Y$ is locally Lipschitz-continuous from $(H\Omega_p(V), \rho) \rightarrow (H\Omega_p(W), \rho)$.*

Proof: Assume $\rho(X, \hat{X}) \leq \epsilon$. This just means that (always for all s, t)

$$|X_{s,t}^k - \hat{X}_{s,t}^k| \leq \epsilon |t - s|^{k/p}, \quad k = 1, 2$$

Introduce an additive control function

$$w(s, t) := (\|X\| \vee \|\hat{X}\| \vee 1)|t - s|.$$

With this choice, the assumptions (4.1) and (4.2) are satisfied and Theorem 4.1 tells us that there exists a constant C , depending on the maximum of the control w and hence only on $\|X\| \vee \|\hat{X}\|$, such that

$$|Y_{s,t}^k - \hat{Y}_{s,t}^k| \leq \epsilon (Cw(s, t))^{k/p}, \quad k = 1, 2.$$

Expanding w we obtain, for a new constant C depending only on $\|X\| \vee \|\hat{X}\|$,

$$\rho(Y, \hat{Y}) = \|Y - \hat{Y}\| \leq \epsilon C$$

The conclusion follows. QED

6. APPLICATION TO THE SUPPORT THEOREM

Consider the Stratonovich-SDE

$$(6.1) \quad dY = f(Y) \circ d\beta$$

with initial condition $Y(0) = y_0 \in W = \mathbb{R}^N$ where $f = (f_1, \dots, f_d)$ stands for d vector-fields, C^3 and globally Lipschitz, driven by

Brownian motion in $V = \mathbb{R}^d$. (The case of an additional drift-term $f_0(Y)dt$ is a trivial generalization of the method below.) In Lyons' form this corresponds to the differential equation

$$dy = f(y)dx$$

lifted to Rough Paths level and using Stratonovich enhanced BM β as driving signal. By Theorem 5.1 the corresponding Itô-map $F : H\Omega_p(V) \rightarrow H\Omega_p(W)$ is continuous w.r.t. the metric ρ . Since $F = (F^1, F^2)$ takes values in $W \oplus W^{\otimes 2}$ there is a natural projection to its underlying path in W . Set

$$\Phi_t(\beta) = y_0 + [F(\beta)]_{0,t}^1.$$

Note that Φ is continuous from $(H\Omega_p(V), \rho)$ to $C^{0,1/p}([0, 1], W)$ with usual Hölder semi-norm (actually norm, since all paths are pinned at y_0 at time 0). Set

$$\Psi(\omega) = \Phi(\beta(\omega)).$$

The map Ψ maps $C([0, 1], V)$ to $C^{0,1/p}([0, 1], W)$ and is measurable only (due to $\omega \rightarrow \beta(\omega)$) and we will also call it Itô-map (no confusion will arise). Note that

$$Y_t(\omega) := \Psi_t(\omega)$$

solves (6.1). Equip $C([0, 1], V)$ with the standard Wiener-measure μ . Our aim is to describe the support of $(\Psi)_*\mu = (\Phi)_*P$ where P is the law of EBM on $H\Omega_p(V)$.

Continuity of the Φ allows to restrict the discussion to the support of P only.

Theorem 6.1. *The support of P equals the ρ -closure of $\mathcal{D}$.*

Proof:

From Corollary 3.4 resp. Corollary 3.5, a.s. and in ρ -metric,

$$\beta^{ad}(n)(\omega) \rightarrow \beta(\omega), \quad \beta(\omega - \omega^{ad}(n) + h) \rightarrow (h^1, h^2),$$

for arbitrary $(h^1, h^2) \in \mathcal{D}$. The first convergence implies that $\text{supp } P$ is contained in the ρ -closure of $\mathcal{D}$. The second convergence result shows, in particular, convergence in probability of $\beta(\omega - \omega^{ad}(n) + h) \rightarrow (h^1, h^2)$. This means that the probability of being within an ϵ -ball from (h^1, h^2) tends to 1, hence is positive for n large enough. By Girsanov's theorem the same is true for $\beta(\omega)$. Hence we get to

other desired inclusion, $\mathcal{D}$ is contained in $\text{supp } P$. (This argument is due to [MS]).

Remark: As in [LQZ] the closure of $\mathcal{D}$ is seen to coincide with the closure of the natural lift of the Cameron-Martin space. This also follows from the results in [FV].

Remarking that solving a rough differential equation driven by a smooth rough path amounts to solve a controlled ODE ([LQ], p164) we have

Corollary 6.2. *The law of the diffusion-process Y in equation (6.1), that is, $(\Psi)_*\mu$ is the $(1/p\text{-H\"older})$ closure of the solutions to the control ODE,*

$$dy = f(y)dh$$

for all $h \in \mathcal{D}$.

(Similarly, one can obtain a support description for the rough path solution to (6.1).)

APPENDIX A. PROOF OF LEMMA (2.2)

Proof. To avoid trivialities assume $s < t$, $s' < t'$. W.l.o.g. $s \leq s'$. Then

$$\begin{aligned} E[|X_{s,t} - X_{s',t'}|^2] &= E[|X_{s,t}|^2] - 2E[X_{s,t}X_{s',t'}] + E[|X_{s',t'}|^2] \\ &= \frac{1}{2}|t-s|^{2-2\alpha} - 2E[X_{s,t}X_{s',t'}] + \frac{1}{2}|t'-s'|^{2-2\alpha}. \end{aligned}$$

To deal with the middle part we distinguish between a few cases.

Case i) $t \leq s'$. By independence of Brownian increments,

$$\begin{aligned} E[|X_{s,t} - X_{s',t'}|^2] &= \frac{1}{2}(|t-s|^{2-2\alpha} + |t'-s'|^{2-2\alpha}) \\ &\leq \frac{1}{2}(|s'-s|^{2-2\alpha} + |t'-t|^{2-2\alpha}) \\ &\leq \left(\frac{|s'-s| + |t'-t|}{2}\right)^{2-2\alpha} \\ &= c(\alpha) \left(|s'-s| + |t'-t|\right)^{2-2\alpha} \end{aligned}$$

where we used $\alpha \in [1/2, 1)$ for the last inequality.

Case ii.1) $s \leq s' < t \leq t'$. Since $X_{s,t} - X_{s',t'} = X_{s,t} - X_{s',t} + X_{s',t} - X_{s',t'}$ we estimate separately $E[|X_{s,t} - X_{s',t}|^2]$ and $E[|X_{s',t} - X_{s',t'}|^2]$. As for the first, the cross-term is readily computed via Ito's isometry and we have

$$E[|X_{s,t} - X_{s',t}|^2] = \frac{1}{2}|t - s|^{2-2\alpha} - \frac{|t - s'|^{2-\alpha}}{|t - s|^\alpha} + \frac{1}{2}|t - s'|^{2-2\alpha}.$$

The r.h.s. is indeed bounded by a constant times $|s' - s|^{2-2\alpha}$ as follows from the lemma below with $a = s' - s, b = t - s'$. As for the second, we get

$$E[|X_{s',t} - X_{s',t'}|^2] = \frac{1}{2}|t - s'|^{2-2\alpha} - \frac{|t' - s'|^{2-\alpha}}{|t - s'|^\alpha} + \frac{1}{2}|t' - s'|^{2-2\alpha}.$$

Choose $a = t' - t, b = t - s'$ and again apply the lemma. Both together yield the required estimate,

$$E[|X_{s,t} - X_{s',t'}|^2] \leq c(|s' - s| + |t' - t|)^{2-2\alpha}.$$

Case ii.2) $s \leq s' \leq t' \leq t$ Similar. QED

Lemma A.1. *For non-negative real numbers $a, b, \alpha \in (0, 1)$ one has*

$$\frac{1}{2}(a + b)^{2-2\alpha} - \frac{b^{2-\alpha}}{(a + b)^\alpha} + \frac{1}{2}b^{2-2\alpha} \leq ca^{2-2\alpha}$$

for some constant $c = c(\alpha)$.

Proof: Divide the l.h.s. by $a^{2-2\alpha}$ and observe that it is a continuous function of $x = b/a \in [0, \infty)$. An easy expansion shows that everything stays bounded as $x \rightarrow \infty$. The lemma follows.

Acknowledgement: Thanks to S.Varadhan, G. Ben Arous and N. Victoir for related discussions, to E. Kosygina for reading an early draft. The author also acknowledges gratefully financial support by the Austrian Academy of Science.

REFERENCES

- [BGL] G. Ben Arous, M. Grădinaru, M. Ledoux, Hölder norms and the support theorem for diffusions. Ann. Inst. H. Poincaré Probab. Statist. 30, no3, 415-436 (1994)
- [FV] P.K.Friz, N.Victoir, forthcoming
- [Gu] M.Gubinelli, Controlling Rough Paths, arXiv:math.PR/0306433 v1
- [IW] IW] N. Ikeda, S. Watanabe, SDEs and diffusion processes. North Holland (1981,89)

- [Le] A. Lejay, Introduction to Rough Paths,
www.iecn.u-nancy.fr/~lejay/rough.html, to appear in Sém. de Proba.,
 LNM, Springer
- [L95] T. Lyons, Interpretation and Solution of ODEs Driven by Rough Signals,
 Proc. Symposia Pure Math. Vol 57, 115-128 (1994)
- [L98] T. Lyons, Differential Equations driven by rough signals, Revista Matem-
 atica Iberoamericana, Vol. 14, No. 2, 215-310 (1998)
- [LQ97] T. Lyons, Z. Qian, Calculus of Variation for Multiplicative Functionals,
 New Trends in Stochastic Analysis, 348-374 Charingworth 1994. World
 Scientific, River Edge, NJ (1997)
- [LQ] T. Lyons, Z. Qian, System Control and Rough Paths, Oxford University
 Press (2002)
- [LQZ] M. Ledoux, Z. Qian, T. Zhang, Large deviations and support theorem
 for diffusion processes via Rough Paths, Stoch. Proc. Appl. 102:2, 265-283
 (2002).
- [M] P. Malliavin, Stochastic Analysis, Springer (1997)
- [MS] A. Millet, M. Sanz-Sole, A simple proof of the support theorem for diffusion
 processes, LNM 1583, pages 36-48
- [RY] D. Revuz, M. Yor, Continuous Martingales and Brownian Motion,
 Springer (1999)
- [S] E. Sippilainen, A pathwise view of solutions to SDEs, unpublished PhD-
 thesis Univ. of Edinburgh (1993)
- [ST] D.W. Stroock, S. Taniguchi, Diffusions as integral curves, or Stratonovich
 without Itô integration, Birkhäuser, Prog. Probab. 34, 333-369 (1994)
- [SV] D.W. Stroock, S.R.S. Varadhan, On the support of diffusion processes with
 application to the strong maximum principle. Proc. 6th Berkeley Symp.
 Math. Statist. Prob. III. Univ. California Press, 333-359 (1972)

COURANT INSTITUTE, NEW YORK UNIVERSITY

E-mail address: `Peter.Friz@cims.nyu.edu`